AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application.

Listing of Claims:

Claim1 (currently amended): A compound of the Formula:

or a pharmaceutically acceptable salt thereof, wherein;

B is phenyl or a heteroaryl of 5 or 6 ring members, wherein a nitrogen with a removable hydrogen or a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to A is optionally substituted by R³², a nitrogen with a removable hydrogen or a carbon at the other position adjacent to the point of attachment is optionally substituted by R³⁶, a nitrogen with a removable hydrogen or a carbon adjacent to R³² and two atoms from the point of attachment is optionally substituted by R³³, a nitrogen with a removable hydrogen or a carbon adjacent to R³⁶ and two atoms from the point of attachment is optionally substituted by R³⁵, and a nitrogen with a removable hydrogen or a carbon adjacent to both R³³ and R³⁵ is optionally substituted by R³⁴;

R⁹, R¹⁰, R¹¹, R¹², R¹³, R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkylenedioxy, haloalkylthio, alkanoyloxy, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, alkoxyalkyl, haloalkoxylalkyl, hydroxy, amino, alkoxyamino, nitro, alkylamino, N-alkyl-N-arylamino, arylamino, aralkylamino, heteroarylamino, heterocyclylamino, heterocyclylamino, alkylthio, alkylthioalkyl, alkylsulfinyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl,

heteroarylsulfinyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, alkylsulfonylalkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkylsulfonamido, amidosulfonyl, alkanoyl, haloalkanoyl, alkyl, alkenyl, halo, haloalkyl, haloalkenyl, haloalkoxy, hydroxyhaloalkyl, hydroxyalkyl, aminoalkyl, haloalkoxyalkyl, carboxyalkyl, carboalkoxy, carboxy, carboxamido, carboxamidoalkyl, and cyano;

B is optionally selected from the group consisting of hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶;

B-is-optionally a C3-C12 cycloalkyl or C4-C9 saturated heterocyclyl, wherein each ring carbon is optionally substituted with R³³, a ring carbon other than the ring carbon at the point of attachment of B to A is optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, ring carbons and a nitrogen adjacent to the carbon atom at the point of attachment are optionally substituted with R⁹ or R⁴³, a ring carbon or nitrogen atom adjacent to the R⁹-position and two atoms from the point of attachment is optionally substituted with R¹⁰, a ring carbon or nitrogen adjacent to the R¹³-position and two atoms from the point of attachment is optionally substituted with R¹², a ring carbon or nitrogen three atoms from the point of attachment and adjacent to the R⁴⁹-position is optionally substituted with R³¹, a ring carbon or nitrogen three atoms from the point of attachment and adjacent to the R³⁴-position is optionally substituted with R³³, and a ring carbon or nitrogen four atoms from the point of attachment and adjacent to the R⁴⁴-and R³⁵-positions is optionally substituted with R³⁴;

A is selected from the group consisting of a bond, $(W^{7})_{rr}$ (CH(R¹⁵))_{pa}, and $(CH(R^{15}))_{pa}$ (W⁷)_{rr} wherein rr is 0 or 1, pa is an integer selected from 0 through 6, and W⁷ is selected from the group consisting of O, S, C(O), (R⁷)NC(O), (R⁷)NC(S), and N(R⁷) with the proviso that no more than one of the group consisting of rr and pa is 0 at the same time;

Ψ is NH or NOH;

M is $[[N or]] R^1-C$;

R¹ is selected from the group consisting of hydrido, alkyl, alkenyl, cyano, halo, haloalkyl, haloalkoxy, haloalkylthio, amino, aminoalkyl, alkylamino, amidino, hydroxy, hydroxyamino, alkoxy, hydroxyalkyl, alkoxyamino, thiol, and alkylthio;

 R^2 is Z^0 -Q:

 Z^0 is selected from the group consisting of a bond, $(CR^{41}R^{42})_q$ -wherein q is an integer selected from 1 through 3, and $(CH(R^{41}))_g$ -W⁰- $(CH(R^{42}))_p$ -wherein g and p are integers independently selected from 0 through 3 and W⁰ is selected from the group consisting of O, S, C(O), S(O), N(R⁴¹), and $ON(R^{41})$;

Z⁰ is optionally (CH(R⁴¹))_e-W²²-(CH(R⁴²))_h wherein e and h are independently 0 or 1 and W²² is selected from the group consisting of CR⁴¹=CR⁴², 1,2-cyclopropyl, 1,2-cyclobutyl, 1,2-cyclohexyl, 1,3-cyclohexyl, 1,2-cyclopentyl, 1,3-cyclopentyl, 2,3-morpholinyl, 2,4-morpholinyl, 2,6-morpholinyl, 3,4-morpholinyl, 3,5-morpholinyl, 1,2-piperazinyl, 1,3-piperazinyl, 2,3-piperazinyl, 2,6-piperazinyl, 1,2-piperidinyl, 1,3-piperidinyl, 2,4-piperidinyl, 2,6-piperidinyl, 3,4-piperidinyl, 1,2-pyrrolidinyl, 1,3-pyrrolidinyl, 2,3-pyrrolidinyl, 2,4-pyrrolidinyl, 2,5-pyrrolidinyl, 3,4-pyrrolidinyl, 2,3-tetrahydrofuranyl, 2,4-tetrahydrofuranyl, 2,5-pyrrolidinyl, 2,5-tetrahydrofuranyl, and 3,4-tetrahydrofuranyl, wherein Z⁰ is directly bonded to the uracil ring and W²² is optionally substituted with one or more substituents selected from the group consisting of R⁹, R¹⁰, R¹¹, R¹², and R¹³;

R⁴¹-and R⁴²-are independently selected from the group consisting of amidino, hydroxyamino, hydrido, hydroxy, amino, and alkyl;

Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein a nitrogen with a removable hydrogen or a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁰, a nitrogen with a removable hydrogen or a carbon at the other position adjacent to the point of

attachment is optionally substituted by R¹³, a nitrogen with a removable hydrogen or a carbon adjacent to R⁹ and two atoms from the point of attachment is optionally substituted by R¹⁰, a nitrogen with a removable hydrogen or a carbon adjacent to R¹³ and two atoms from the point of attachment is optionally substituted by R12, and a nitrogen with a removable hydrogen or a carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R11; wherein (a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², and (e) a ring carbon, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹;

Q is optionally hydrido with the proviso that Z⁰ is selected from other than a bond;

K is (CR^{4a}R^{4b})_n wherein n is 1 or 2;

R^{4a} and R^{4b} are independently selected from the group consisting of halo, hydrido, hydroxyalkyl, alkyl, alkoxyalkyl, alkylthioalkyl, and haloalkyl;

 E^0 is E^1 , when K is $(CR^{4a}R^{4b})_n$, wherein E^1 is selected from the group consisting of a bond, C(O), C(S), C(O)N(R⁷), (R⁷)NC(O), S(O)₂, (R⁷)NS(O)₂, and S(O)₂N(R⁷);

Y⁰ is

phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q^s, a carbon two or three atoms from the point of attachment of Q^s to said phenyl or said heteroaryl to said phenyl or said heteroaryl is substituted by Q^b, a carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁷, another carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁸, a carbon adjacent to Q^b is optionally substituted by R¹⁶, and another carbon adjacent to Q^b is optionally substituted by R¹⁹;

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, nitro, alkoxyamino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, alkenyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, haloalkoxyalkyl, carboalkoxy, and cyano[[;]].

R¹⁶-or R¹⁹ is optionally selected from the group consisting of NR²⁰R²¹, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁶, R¹⁹, and Q^b-are not simultaneously hydrido;

Q^b-is selected from the group consisting of NR²⁰R²¹, aminoalkyl, hydrido, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the provise that no more than one of R²⁰ and R²¹ is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino at the same time, with the further provise that no more than one of R²³ and R²⁴ is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino at the same time;

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, alkyl, hydroxy, aminoalkyl, amino, dialkylamino, alkylamino, and hydroxyalkyl;

 Q^6 -is selected from the group consisting of bond, $(CR^{37}R^{38})_b$ wherein b is an integer selected from 1 through 4, and $(CH(R^{14}))_c$ -W¹- $(CH(R^{15}))_d$ wherein c and d are integers independently selected from 1 through 3 and W¹-is selected from the group consisting of $C(O)N(R^{14})$, $(R^{14})NC(O)$, S(O), S(O)₂, S(O)₂ $N(R^{14})$, $N(R^{14})S(O)$ ₂, and $N(R^{14})$, with the provise that R^{14} -is selected from other than halo when directly bended to N, with the further provisen that Q^6 -is selected from other than a bend when Y^0 -is $Z-Q^b$ -5- Q^6 -6- R^{17} -4- R^{18} -3- R^{19} -pyridine or $Z-Q^b$ -4- Q^6 -3- R^{16} -5- R^{18} -6- R^{19} -pyridine, and with the additional provise that $(CR^{37}R^{38})_b$ and $(CH(R^{14}))_c$ are bended to E^0 :

R¹⁴-is-selected from the group consisting of hydrido, halo, alkyl, and haloalkyl; R³⁷-and R³⁸-are independently selected from the group consisting of hydrido, alkyl, and haloalkyl;

R³⁸ is optionally aroyl or heteroaroyl, wherein R³⁸ is optionally substituted with one or more substituents selected from the group consisting of R¹⁶, R¹⁷, R¹⁸, and R¹⁹;

Y⁰-is optionally Y^{AT}-wherein Y^{AT}-is-Q^b-Q^s;

 Y^0 is optionally Q^b - Q^{ss} wherein Q^{ss} is $(CH(R^{14}))_e$ - W^2 - $(CH(R^{15}))_h$, wherein e and h are independently 1 or 2 and W^2 is CR^{4a} = CR^{4b} , with the proviso that $(CH(R^{14}))_e$ is bonded to E^0 ;

Y⁰-is-optionally Q^b-Q^{sess}-or Q^b-Q^{sessr}-wherein Q^{sess}-is (CH(R³⁸))_r-W⁵, Q^{sessr}-is (CH(R³⁸))_r-W⁶, r is 1 or 2, W⁵-and W⁶-are independently selected from the group consisting of 1,4-indenyl, 1,5-indenyl, 1,6-indenyl,

- 1,7-indenyl, 2,7-indenyl, 2,6-indenyl, 2,5-indenyl, 2,4-indenyl, 3,4-indenyl,
- 3,5-indenyl, 3,6-indenyl, 3,7-indenyl, 2,4-benzofuranyl, 2,5-benzofuranyl,
- 2,6-benzofuranyl, 2,7-benzofuranyl, 3,4-benzofuranyl, 3,5-benzofuranyl,
- 3,6-benzofuranyl, 3,7-benzofuranyl, 2,4-benzothiophenyl, 2,5-benzothiophenyl, 2,6-benzothiophenyl, 2,7-benzothiophenyl, 3,4-benzothiophenyl,
- 3,5-benzothiophenyl, 3,6-benzothiophenyl, 3,7-benzothiophenyl,
- 2,7-imidazo(1,2-a)pyridinyl, 3,4-imidazo(1,2-a)pyridinyl,
- 3,5-imidazo(1,2-a)pyridinyl, 3,6-imidazo(1,2-a)pyridinyl,

3,7-imidazo(1,2-a)pyridinyl, 2,4-indolyl, 2,5-indolyl, 2,6-indolyl, 2,7-indolyl, 3,4indolyl, 3,5-indolyl, 3,6-indolyl, 3,7-indolyl, 1,4-isoindolyl, 1,5-isoindolyl, 1,6isoindolyl, 2,4-isoindolyl, 2,5-isoindolyl, 2,6-isoindolyl, 2,7-isoindolyl, 1,3-isoindolyl, 3,4-indazolyl, 3,5-indazolyl, 3,6-indazolyl, 3,7-indazolyl, 2,4-benzoxazolyl, 2,5-benzoxazolyl, 2,6-benzoxazolyl, 2,7-benzoxazolyl, 3,4-benzisoxazolyl, 3,5-benzisoxazolyl, 3,6-benzisoxazolyl, 3,7-benzisoxazolyl, 1,4naphthyl, 1,5-naphthyl, 1,6-naphthyl, 1,7-naphthyl, 1,8-naphthyl, 2,4-naphthyl, 2,5-naphthyl, 2,6-naphthyl, 2,7-naphthyl, 2,8-naphthyl, 2,4-quinolinyl, 2,5-quinolinyl, 2,6-quinolinyl, 2,7-quinolinyl, 2,8-quinolinyl, 3,4-quinolinyl, 3,5-quinolinyl, 3,6-quinolinyl, 3,7-quinolinyl, 3,8-quinolinyl, 4,5-quinolinyl, 4,6-quinolinyl, 4,7-quinolinyl, 4,8-quinolinyl, 1,4-isoquinolinyl, 1,5isoquinolinyl, 1,6-isoquinolinyl, 1,7-isoquinolinyl, 1,8-isoquinolinyl, 3,4-isoquinolinyl, 3,5-isoquinolinyl, 3,6-isoquinolinyl, 3,7-isoquinolinyl, 3,8-isoquinolinyl, 4,5-isoquinolinyl, 4,6-isoquinolinyl, 4,7-isoquinolinyl, 4,8-isoquinolinyl, 3,4-cinnolinyl, 3,5-cinnolinyl, 3,6-cinnolinyl, 3,7-cinnolinyl, 3,8cinnolinyl, 4,5-cinnolinyl, 4,6-cinnolinyl, 4,7-cinnolinyl, and 4,8-cinnolinyl, and each carbon and hyrido containing nitrogen member of the ring of the W⁵ and of the ring of the W⁶, other than the points of attachment of W⁵ and W⁶, is optionally substituted with one or more of the group consisting of R⁹, R¹⁰, R¹¹, and R¹², with the proviso that Q^b is bonded to lowest number substituent position of each W⁵, with the further proviso that Q^b is bonded to highest number substituent position of each W⁶, and with the additional proviso that (CH(R³⁸))_r is bonded to E⁰.

Claim 2 (currently amended): Compound of [[Claim]] claim 1 of the Formula:

or a pharmaceutically acceptable salt thereof, wherein;

B is phenyl or a heteroaryl of 5 or 6 ring members, wherein a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to A is optionally substituted by R³², the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R³⁶, a carbon adjacent to R³² and two atoms from the carbon at the point of attachment is optionally substituted by R³³, a carbon adjacent to R³⁶ and two atoms from the carbon at the point of attachment is optionally substituted by R³⁵, and any carbon adjacent to both R³³ and R³⁵ is optionally substituted by R³⁴;

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkylenedioxy, haloalkylthio, alkanoyloxy, alkoxy, hydroxy, amino, alkoxyamino, haloalkanoyl, nitro, alkylamino, alkylthio, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkylsulfonamido, amidosulfonyl, alkyl, alkenyl, halo, haloalkyl, haloalkenyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b;

B is optionally selected from the group consisting of hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶;

B is optionally a C3-C12 cycloalkyl or a C4-C9 saturated heterocyclyl, wherein each ring carbon is optionally substituted with R³³, a ring carbon other than the ring carbon at the point of attachment of B to A is optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, ring carbons and a nitrogen adjacent to the carbon atom at the point of attachment are optionally substituted with R⁹ or R¹³, a ring carbon or nitrogen atom adjacent to the R⁹ position and two atoms from the point of attachment is optionally substituted with R¹⁰, a ring carbon or nitrogen atom adjacent to the R¹³ position and two atoms from the point of attachment is optionally substituted with R¹², a ring carbon or nitrogen atom three atoms from the point of attachment and adjacent to the R¹⁰ position is optionally substituted with R¹¹, a ring carbon or nitrogen atom three atoms from the point of attachment and adjacent to the R¹² position is optionally substituted with R³³, and a ring carbon or nitrogen atom four atoms from the point of attachment and adjacent to the R³⁴;

R⁹, R¹⁰, R¹¹, R¹², and R¹³ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, alkoxyamino, alkanoyl, haloalkanoyl, amidino, guanidino, alkylenedioxy, haloalkylthio, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, hydroxy, amino, alkylamino, N-alkyl-N-arylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, heterocyclylamino, heterocyclylalkylamino, alkylthio, alkylsulfinyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, alkylsulfamido, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, amidosulfonyl, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, and cyano;

A is a bond or $(CH(R^{15}))_{pa}$ - $(W^7)_{rr}$ wherein rr is 0 or 1, pa is an integer selected from 0 through 3, and W^7 is selected from the group consisting of O, S, C(O), $(R^7)NC(O)$, $(R^7)NC(S)$, and $N(R^7)$;———

R⁷ is selected from the group consisting of hydrido, hydroxy and alkyl;

——R¹⁵-is selected from the group consisting of hydrido, hydroxy, halo, alkyl, and haloalkyl;

M-is-R¹-C:

R¹ is selected from the group consisting of hydrido, alkyl, cyano, halo, haloalkyl, haloalkoxy, amino, aminoalkyl, alkylamino, amidino, hydroxy, hydroxyamino, alkoxy, hydroxyalkyl, alkoxyamino, thiol, and alkylthio;

 R^2 is Z^0 -Q:

 Z^{0} is selected from the group consisting of a bond, $(CR^{41}R^{42})_{q}$ wherein q is 1 or 2, and $(CH(R^{41}))_{g}$ W^{0} - $(CH(R^{42}))_{p}$ wherein g and p are integers independently selected from 0 through 3 and W^{0} is selected from the group consisting of O, S, C(O), $N(R^{41})$, and $ON(R^{41})$:

Z⁰-is optionally (CH(R⁴¹))_e-W²²-(CH(R⁴²))_h-wherein e and h are independently 0 or 1 and W²²-is selected from the group consisting of CR⁴¹=CR⁴², 1,2-cyclopropyl, 1,2-cyclobutyl, 1,2-cyclohexyl, 1,3-cyclohexyl, 1,2-cyclopentyl, 1,3-cyclopentyl, 2,3-morpholinyl, 2,4-morpholinyl, 2,6-morpholinyl, 3,4-morpholinyl, 3,5-morpholinyl, 1,2-piperazinyl, 1,3-piperazinyl, 2,3-piperazinyl, 2,6-piperazinyl, 1,2-piperidinyl, 1,3-piperidinyl, 2,4-piperidinyl, 2,6-piperidinyl, 3,4-piperidinyl, 1,2-pyrrolidinyl, 2,3-pyrrolidinyl, 2,3-pyrrolidinyl, 2,4-pyrrolidinyl, 2,4-pyrrolidinyl, 2,5-pyrrolidinyl, 3,4-pyrrolidinyl, 2,3-tetrahydrofuranyl, 2,4-tetrahydrofuranyl, 2,5-tetrahydrofuranyl, and 3,4-tetrahydrofuranyl, wherein Z⁰-is directly bonded to the uracil ring and W²²-is optionally substituted with one or more substituents selected from the group consisting of R⁹, R¹⁰, R¹¹, R¹², and R¹³;

R⁴¹ and R⁴² are independently selected from the group consisting of hydrido, hydroxy, alkyl, and amino;

Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁰, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁰ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹;

Q is optionally hydrido with the proviso that Z⁰ is other than a bond;

K is CHR^{4a} wherein R^{4a} is selected from the group consisting of hydrido, hydroxyalkyl, alkyl, alkoxyalkyl, alkylthioalkyl, and haloalkyl;

E⁰ is selected from the group consisting of a covalent single bond, C(O)N(H), (H)NC(O), (R⁷)NS(O)₂, and S(O)₂N(R⁷);

Y⁰-is phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q⁶, a carbon two or three-atoms from the point of attachment of Q⁶-to said phenyl or said heteroaryl is substituted by Q⁶, a carbon adjacent to the point of attachment of Q⁶ is optionally substituted by R¹⁷, another carbon adjacent to the point of attachment of Q⁶-is optionally substituted by R¹⁸, a carbon adjacent to Q⁶-is optionally substituted by R¹⁶, and another carbon adjacent to Q⁶-is optionally substituted by R¹⁹;

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano[[;]].

R¹⁶-or R¹⁹-is optionally selected from the group consisting of NR²⁰R²¹, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁶, R¹⁹, and Q^b-are not simultaneously hydrido;

Q^b-is selected from the group consisting of NR²⁰R²¹, hydrido, N(R²⁵)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that no more than one of R²⁰-and R²¹-is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino at the same time, with the further proviso that no more than one of R²³-and R²⁴-is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino at the same time;

——R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, alkyl, hydroxy, amino, alkylamino and dialkylamino;

Q⁵ is selected from the group consisting of a bond, (CR³⁷R³⁸)_b wherein b is an integer selected from 1 through 4, and (CH(R¹⁴))_e-W¹-(CH(R¹⁵))_d wherein c and d are integers independently selected from 1 through 3 and W¹ is selected from the group consisting of C(O)N(R¹⁴), (R¹⁴)NC(O),

S(O), S(O)₂, S(O)₂N(R¹⁴), N(R¹⁴)S(O)₂, and N(R¹⁴), with the proviso that R¹⁴ is selected from other than halo when directly bonded to N and with the further proviso that (CR³⁷R³⁸)_b, and (CH(R¹⁴))_c are bonded to E⁰;

R¹⁴-is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl; R³⁷-and R³⁸-are independently selected from the group consisting of hydrido, alkyl, and haloalkyl;

R³⁸ is optionally aroyl or heteroaroyl, wherein R³⁸ is optionally substituted with one or more substituents selected from the group consisting of R¹⁶, R¹⁷, R¹⁸, and R¹⁹:

Y⁰ is optionally Y^{AT} wherein Y^{AT} is Q^b-Q⁵;

 Y^0 -is-optionally Q^b - Q^{sc} -wherein Q^{sc} -is $(CH(R^{14}))_e$ - W^2 - $(CH(R^{15}))_h$, wherein e and h are integers independently selected from 1 through 2 and W^2 -is CR^{4a} =CH with the proviso that $(CH(R^{14}))_e$ is bonded to E^0 .

Claims 3-16 (cancelled)

Claim 17(currently amended): Compound of [[Claim]] claim 2 of the Formula:

$$\begin{array}{c|c} & & & \\ &$$

or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, C2-C8 alkyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶;

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b;

A is a bond or $(CH(R^{15}))_{pa}$ - $(W^7)_{rr}$ wherein rr is 0 or 1, pa is an integer selected from 0 through 3, and W^7 is $(R^7)NC(O)$ or $N(R^7)$;

R⁷-is-selected from the group consisting of hydrido, hydroxy and alkyl;

R¹⁵-is-selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

M is [[N or]] R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxyamino, amidino, amino, cyano, hydroxyalkyl, alkoxy, alkyl, alkylamino, aminoalkyl, alkylthio, alkoxyamino, haloalkyl, haloalkoxy, and halo;

 R^2 is Z^0 -Q:

 Z^0 is selected from the group consisting of a bond, CH_2 , CH_2 , CH_2 , W^0 - $(CH(R^{42}))_p$ wherein p is 0 or 1 and W^0 is selected from the group consisting of O, S, and $N(R^{41})$;

R⁴¹ and R⁴² are independently hydrido or alkyl;

Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹; wherein (a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring

carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11} ;

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkylsulfonamido, alkylsulfinyl, alkylsulfonyl, amidosulfonyl, alkyl, alkoxy, halo, haloalkyl, hydroxyhaloalkyl, carboxy, carboxamido, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy,heterocyclyloxy, heterocyclylalkoxy, hydroxy, amino, alkoxyamino, alkylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, heterocyclylamino, heterocyclylalkylamino, alkylsulfonamido, amidosulfonyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, arylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, halo, haloalkyl, and cyano;

$$R^{17}$$
 R^{16}
 NH
 NH_2
 R^{18}
 R^{19} ; and

phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q⁶, a carbon two or three atoms from the point of attachment of Q⁶ to said phenyl or said heteroaryl is substituted by Q⁶, a carbon adjacent to the point of attachment of Q⁶ is optionally substituted by R¹⁷, another carbon

adjacent to the point of attachment of Q⁵ is optionally substituted by R¹⁸, a carbon adjacent to Q^b is optionally substituted by R¹⁶, and another carbon adjacent to Q^b is optionally substituted by R¹⁹;

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, alkylamino, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano[[;]].

R¹⁶-or R¹⁹-is optionally selected from the group consisting of NR²⁰R²¹, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;

Q^b-is selected from the group consisting of NR²⁰R²¹, hydrido,

C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), with the proviso that no more than one of R²⁰ and R²¹-is hydroxy at the same time and with the further proviso that no more than one of R²³ and R²⁴-is hydroxy at the same time;

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, alkyl, and hydroxy;

Q^s is selected from the group consisting of a bond, CH₂, and CH₂CH₂.

Claim 18 (currently amended): Compound of [[Claim]] claim 17 or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, ethyl, 2-propynyl, 2-propenyl, propyl, isopropyl, butyl, 2-butenyl, 3-butenyl, 2-butynyl, sec-butyl, tert-butyl, isobutyl, 2-methylpropenyl, 1-pentyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 2-pentynyl, 3-pentynyl, 2-pentyl, 1-methyl-2-butenyl, 1-methyl-3-butenyl, 1-methyl-2-butynyl, 3-pentyl, 1-ethyl-2-propenyl, 2-methylbutyl, 2-methyl-2-butenyl, 2-methyl-3-butenyl, 2-methyl-3-butynyl, 3-methylbutyl, 3-methyl-2-butenyl, 3-methyl-3-butenyl, 1-hexyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 2-hexyl, 1-methyl-2-pentenyl, 1-methyl-3-pentenyl, 1-methyl-4-pentenyl, 1-methyl-2-butenyl, 1-methyl-3-pentynyl, 3-hexyl, 1-ethyl-2-butenyl,

1-ethyl-3-butenyl, 1-propyl-2-propenyl, 1-ethyl-2-butynyl, 1-heptyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 6-heptenyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 5-heptynyl, 1-methyl-2-hexenyl, 1-methyl-3-hexenyl, 1-methyl-4-hexenyl, 1-methyl-2-hexynyl, 1-methyl-3-hexynyl, 1-methyl-4-hexynyl, 3-heptyl, 1-ethyl-2-pentenyl, 1-ethyl-3-pentenyl, 1-ethyl-4-pentenyl, 1-butyl-2-propenyl, 1-ethyl-2-pentynyl, 1-ethyl-3-pentynyl, 2,2,2-trifluoroethyl, 2,2-difluoropropyl, 4-trifluoromethyl-5,5,5-trifluoropropyl, 4-trifluoromethylpentyl, 5,5,6,6,6-pentafluorohexyl, and 3,3,3-trifluoropropyl, wherein each member of group B is optionally substituted at any carbon up to and including 5 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶;

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, amidosulfonyl, N-methylamidosulfonyl, N,N-dimethylamidosulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, cyano, and Q^b;

A is selected from the group consisting of bond, NH, N(CH₃), N(OH), CH₂, CH₃CH, CF₃CH, NHC(O), N(CH₃)C(O), C(O)NH, C(O)N(CH₃), CH₂CH₂, CH₂CH₂, and CF₃CHCH₂;

A is optionally selected from the group consisting of CH₂N(CH₃), CH₂CH₂N(CH₃), and CH₂CH₂N(CH₂CH₃) with the proviso that B is hydrido;

-----M is N or R¹-C:

R¹ is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, 1-aminoethyl, methylamino, dimethylamino, cyano, methyl,

ethyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, methoxy, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxyamino, methylthio, ethylthio, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, and bromo;

Z⁰ is selected from the group consisting of a bond, CH₂, CH₂CH₂, O, S, NH, N(CH₃), OCH₂, SCH₂, N(H)CH₂, and N(CH₃)CH₂;——

Q is selected from the group consisting of phenyl, 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 3-isoxazolyl, 5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, and 1,3,5-triazin-2-yl, wherein a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁴, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁴, and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹;

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, N-methylamino, N,N-dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, methanesulfonamido, amidosulfonyl, N-methylamidosulfonyl, N,N-dimethylamidosulfonyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, carboxymethyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, aminomethyl,

1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methanesulfonamido, amidosulfonyl, N-methylamidosulfonyl,

N,N-dimethylamidosulfonyl, hydroxymethyl, 1-hydroxyethyl,

2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl,

N,N-dimethylamidocarbonyl, N-benzylamidocarbonyl,

N-(2-chlorobenzyl)amidocarbonyl, N-(3-fluorobenzyl)amidocarbonyl,

N-(2-trifluoromethylbenzyl)amidocarbonyl, N-(1-phenylethyl)amidocarbonyl, N-(1-methyl-1-phenylethyl)amidocarbonyl, N-benzylamidosulfonyl,

N-(2-chlorobenzyl)amidosulfonyl, N-ethylamidocarbonyl,

N-isopropylamidocarbonyl, N-propylamidocarbonyl,

N-isobutylamidocarbonyl, N-(2-butyl)amidocarbonyl,

N-cyclobutylamidocarbonyl, N-cyclopentylamidocarbonyl,

N-cyclohexylamidocarbonyl, fluoro, chloro, bromo, cyano, cyclobutoxy, cyclohexoxy, cyclohexylmethoxy, 4-trifluoromethycyclohexylmethoxy, cyclopentoxy, benzyl, benzyloxy, 4-bromo-3-fluorophenoxy,

3-bromobenzyloxy, 4-bromobenzyloxy, 4-bromobenzylamino,

5-bromopyrid-2-ylmethylamino, 4-butoxyphenamino, 3-chlorobenzyl,

4-chlorophenoxy, 4-chloro-3-ethylphenoxy, 4-chloro-3-ethylbenzylamino,

4-chloro-3-ethylphenylamino, 3-chlorobenzyloxy, 4-chlorobenzyloxy,

4-chlorobenzylsulfonyl, 4-chlorophenylamino, 4-chlorophenylsulfonyl,

5-chloropyrid-3-yloxy, 2-cyanopyrid-3-yloxy, 2,3-difluorobenzyloxy,

2,4-difluorobenzyloxy, 3,4-difluorobenzyloxy, 2,5-difluorobenzyloxy,

3,5-difluorophenoxy, 3,5-difluorobenzyloxy, 4-difluoromethoxybenzyloxy,

2,3-difluorophenoxy, 2,4-difluorophenoxy, 2,5-difluorophenoxy,

3,5-dimethylphenoxy, 3,4-dimethylphenoxy, 3,4-dimethylbenzyloxy,

3,5-dimethylbenzyloxy, 4-ethoxyphenoxy, 4-ethylbenzyloxy, 3-ethylphenoxy, 4-ethylaminophenoxy, 3-ethyl-5-methylphenoxy, 4-fluorobenzyloxy,

2-fluoro-3-trifluoromethylbenzyloxy, 3-fluoro-5-trifluoromethylbenzyloxy,

4-fluoro-2-trifluoromethylbenzyloxy, 4-fluoro-3-trifluoromethylbenzyloxy,

2-fluorophenoxy, 4-fluorophenoxy, 2-fluoro-3-trifluoromethylphenoxy,

2-fluorobenzyloxy, 4-fluorophenylamino, 2-fluoro-4-trifluoromethylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy,

4-isopropyl-3-methylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy,

4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, phenylamino,

1-phenylethoxy, 2-phenylethoxy, 2-phenylethyl, 2-phenylethylamino, phenylsulfonyl,

3-trifluoromethoxybenzyloxy, 4-trifluoromethoxybenzyloxy,

3-trifluoromethoxyphenoxy, 4-trifluoromethoxyphenoxy,

3-trifluoromethylbenzyloxy, 4-trifluoromethylbenzyloxy,

2,4-bis-trifluoromethylbenzyloxy, 3-trifluoromethylbenzyl,

3,5-bis-trifluoromethylbenzyloxy, 4-trifluoromethylphenoxy,

3-trifluoromethylphenoxy, 3-trifluoromethylthiobenzyloxy,

4-trifluoromethylthiobenzyloxy, 2,3,4-trifluorophenoxy, 2,3,5-trifluorophenoxy, 3-pentafluoroethylphenoxy, 3-(1,1,2,2-tetrafluoroethoxy)phenoxy, and

3-trifluoromethylthiophenoxy; and

Y⁰ is selected from the group consisting of:

1-Q^b-4-Q⁶-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹benzene,

2-Q^b-5-Q⁶-6-R¹⁷-4-R¹⁸-3-R¹⁹pyridine,

3-Q^b-6-Q^c-2-R¹⁶-5-R¹⁸-4-R¹⁹pyridine, 2-Q^b-5-Q^c-3-R¹⁶-6-R¹⁸pyrazine, 3-Q^b-6-Q^c-2-R¹⁸-5-R¹⁸-4-R¹⁹pyridazine,

2-Q^b-5-Q^c-4-R¹⁷-6-R¹⁸pyrimidine, 5-Q^b-2-Q^c-4-R¹⁶-6-R¹⁹pyrimidine,

3-Q^b-5-Q^c-4-R¹⁶-2-R¹⁹thiophene, 2-Q^b-5-Q^c-3-R¹⁶-4-R¹⁷thiophene,

3-Q^b-5-Q^c-4-R¹⁶-2-R¹⁹furan, 2-Q^b-5-Q^c-3-R¹⁶-4-R¹⁷furan,

3-Q^b-5-Q^c-4-R¹⁶-2-R¹⁹pyrrole, 2-Q^b-5-Q^c-3-R¹⁶-4-R¹⁷pyrrole,

4-Q^b-2-Q^c-5-R¹⁹imidazole, 2-Q^b-4-Q^c-5-R¹⁷imidazole,

3-Q^b-5-Q^c-4-R¹⁶isoxazole, 5-Q^b-3-Q^c-4-R¹⁶isoxazole,

2-Q^b-5-Q^c-4-R¹⁶pyrazole, 4-Q^b-2-Q^c-5-R¹⁹thiazole, and

2-Q^b-5-Q⁶-4-R¹⁷thiazole;

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, methyl, ethyl, isopropyl, propyl, carboxy, amidino, guanidino, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio,

isopropylthio, trifluoromethylthio, methylsulfinyl, ethylsulfinyl, methylsulfonyl, ethylsulfonyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, and cyano[[;]].

R¹⁶-or R¹⁹-is optionally selected from the group-consisting of NR²⁰R²¹, C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;

— Q^b is selected from the group consisting of NR²⁰R²¹, hydrido, C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), with the proviso that no more than one of R²⁰ and R²¹ is hydroxy at the same time and with the further proviso that no more than one of R²³ and R²⁴ is hydroxy at the same time;

—— R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, methyl, ethyl, propyl, butyl, isopropyl, and hydroxy;

—— Q⁶ is selected from the group consisting of a bond, CH₂, and CH₂CH₂.

Claim19 (currently amended): Compound of [[Claim]] claim 18 or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butyl, (R)-2-butyl, (S)-2-butyl, *tert*-butyl, isobutyl, 1-pentyl, 3-pentyl, 2-methylbutyl, 2,2,2-trifluoroethyl, 6-amidocarbonylhexyl, 4-methyl-2-pentyl, 3-hydroxypropyl, 1-methoxy-2-propyl, 2-methoxyethyl, 2-methyl-2-butyl, 3-methyl-2-butyl, 2-dimethylaminopropyl, 2-cyanoethyl, 6-hydroxyhexyl, 2-hydroxyethyl, 2-amidinoethyl, 2-guanidinoethyl, 3-guanidinopropyl, 4-guanidinobutyl, 3-hydroxypropyl, 4-hydroxybutyl, 6-cyanohexyl, 2-dimethylaminoethyl, 3-methylbutyl, 2-methylbutyl, (S)-2-methylbutyl, 3-aminopropyl, 2-hexyl, and 4-aminobutyl;

A is selected from the group consisting of a bond, CH₂, NHC(O), CH₂CH₂, CH₂CH₂, and CH₃CHCH₂;

— M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, methylamino, cyano, methyl, trifluoromethyl, methoxy, hydroxymethyl, methoxyamino, methylthio, trifluoromethoxy, fluoro, and chloro;

Z⁰ is selected from the group consisting of a bond, CH₂, O, S, NH, N(CH₃), OCH₂, and SCH₂;

Q is selected from the group consisting of

3-amidocarbonyl-5-aminophenyl, 3-amino-5-(N-benzylamidocarbonyl)phenyl, 3-amino-5-benzylphenyl, 3-amino-5-(2-phenylethyl)phenyl,

3-amino-5-benzylaminophenyl, 3-amino-5-(2-phenylethylamino)phenyl,

3-amino-5-benzyloxyphenyl, 3-amino-5-(2-phenylethoxy)phenyl,

3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl,

3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl,

3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl,

3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl,

3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl,

3-amino-5-(N-benzylamidosulfonyl)phenyl,

3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl,

3-amino-5-(N-ethylamidocarbonyl)phenyl,

3-amino-5-(N-isopropylamidocarbonyl)phenyl,

3-amino-5-(N-propylamidocarbonyl)phenyl,

3-amino-5-(N-isobutylamidocarbonyl)phenyl,

3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl,

3-amino-5-(N-cyclobutylamidocarbonyl)phenyl,

3-amino-5-(N-cyclopentylamidocarbonyl)phenyl,

3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 5-amino-2-fluorophenyl,

3-amino-5-hydroxymethylphenyl, 5-amino-3-methoxycarbonylphenyl,

3-amidinophenyl, 3-amino-2-methylphenyl, 5-amino-2-methylthiophenyl,

3-aminophenyl, 3-amino-5-(4-trifluoromethylbenzylamino)phenyl,

3-amino-5-(4-trifluoromethylbenzyloxy)phenyl, 3-carboxyphenyl,

3-carboxy-5-hydroxyphenyl, 3-amino-5-carboxyphenyl, 3-chlorophenyl,

2-chlorophenyl, 3-cyanophenyl, 3,5-diaminophenyl, 3-dimethylaminophenyl,

2-fluorophenyl, 3-fluorophenyl, 2-hydroxyphenyl, 3-hydroxyphenyl,

3-methanesulfonylaminophenyl, 2-methoxyphenyl, 3-methoxyphenyl,

3-methoxyaminophenyl, 3-methoxycarbonylphenyl, 2-methylaminophenyl,

3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, phenyl, 3-trifluoroacetamidophenyl, 3-trifluoromethylphenyl, and

2-trifluoromethylphenyl, 5-amino-2-thienyl, 5-amino-3-thienyl,

3-bromo-2-thienyl, 3-pyridyl, 4-pyridyl, 2-thienyl, and 3-thienyl;

Y⁰ is selected from the group consisting of:

1-Q^b-4-Q^c-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹benzene.

2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹pyridine,

3-Q^b-6-Q^c-2-R¹⁶-5-R¹⁸-4-R¹⁹pyridine,

3-Q^b-5-Q^c-4-R¹⁶-2-R¹⁹thiophene, and 2-Q^b-5-Q^c-3-R¹⁶-4-R¹⁷thiophene;

R¹⁶ and R¹⁹ are independently selected from the group consisting of hydrido, amidino, amino, aminomethyl, methoxy, methylamino, hydroxy, hydroxymethyl, fluoro, chloro, and cyano; <u>and</u>

R¹⁶-or R¹⁹-is-optionally C(NR²⁵)NR²³R²⁴-with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrido, fluoro, chloro, hydroxy, hydroxymethyl, amino, carboxy, and cyano[[;]].

Q^b is C(NR²⁵)NR²³R²⁴ or hydrido;

———R²³, R²⁴, and R²⁵ are independently hydrido or methyl;

——Q⁶ is CH₂.

Claim 20 (currently amended): Compound of Claim 17, of the Formula:

$$\begin{array}{c|c} & & & \\ & & \\ & & & \\ & & & \\ & & & \\ & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ &$$

or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, C2-C8 alkyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶;

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b;

A is a bond or $(CH(R^{15}))_{pa}$ $(W^7)_{rr}$ wherein rr is 0 or 1, pa is an integer selected from 0 through 3, and W^7 is $N(R^7)$;

R² is hydrido or alkyl;

———R¹⁵ is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;
———M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxyamino, amidino, amino, cyano, hydroxyalkyl, alkoxy, alkyl, alkylamino, aminoalkyl, alkylthio, alkoxyamino, haloalkyl, haloalkoxy, and halo;

- Z⁰ is a bond:

Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁰, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁰ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹;

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkoxy, alkylsulfinyl, alkylsulfonyl, amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, carboxy, carboxamido, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, alkoxy, alkoxyamino, hydroxy, amino, alkylamino, alkylsulfonamido, amidosulfonyl, hydroxyalkyl, aminoalkyl, halo, haloalkyl, carboalkoxy, carboxy, carboxamido, carboxyalkyl, and cyano; and

Y⁰-is phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q⁶, a carbon two or three atoms from the point of attachment of Q⁶-to said phenyl or said heteroaryl is substituted by Q⁶, a carbon adjacent to the point of attachment of Q⁶ is optionally substituted by R¹⁷, another carbon adjacent to the point of attachment of Q⁶-is optionally substituted by R¹⁸, a carbon adjacent to Q⁶-is optionally substituted by R¹⁶, and another carbon adjacent to Q⁶-is optionally substituted by R¹⁶;

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, alkylamino, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano[[;]].

R¹⁶-or-R¹⁹-is optionally selected from the group consisting of NR²⁰R²¹, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁶, R¹⁹, and Q^b-are not simultaneously hydrido;

Claim 21 (currently amended): Compound of [[Claim]] claim 20 or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butenyl, 2-butynyl, sec-butyl, *tert*-butyl, isobutyl, 2-methylpropenyl, 1-pentyl, 2-pentenyl, 3-pentenyl, 2-pentynyl, 3-pentynyl, 2-pentyl, 3-pentyl, 2-methylbutyl, 2-methyl-2-butenyl, 3-methylbutyl, 3-methyl-2-butenyl, 1-hexyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 2-hexyl, 1-methyl-3-pentynyl, 3-hexyl, 1-methyl-3-pentyl, 1-methyl-2-pentyl, 1-methyl-3-pentyl, 5-heptenyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 5-heptynyl, 2-heptyl, 1-methyl-2-hexenyl, 1-methyl-3-hexenyl, 1-methyl-4-hexenyl, 1-methyl-2-pentenyl, 1-methyl-3-pentenyl, 1-methyl-4-hexynyl, 3-heptyl, 1-ethyl-2-pentenyl, 1-ethyl-3-pentenyl, 1-ethyl-3-pentynyl, 2,2,2-trifluoroethyl, 2,2-difluoropropyl, 4-trifluoromethyl-5,5,5-trifluoropentyl,

4-trifluoromethylpentyl, 5,5,6,6,6-pentafluorohexyl, and 3,3,3-trifluoropropyl, wherein each member of group B is optionally substituted at any carbon up to and including 5 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶;

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, amidino, guanidino, methyl, ethyl, methoxy, ethoxy, hydroxy, amino, N-methylamino, dimethylamino, methoxyamino, methylthio, ethylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, fluoro, chloro, bromo, amidosulfonyl, N-methylamidosulfonyl, hydroxymethyl, amidocarbonyl, carboxy, cyano, and Q^b;

A is selected from the group consisting of a bond, NH, N(CH₃), CH₂, CH₃CH, and CH₂CH₂;

A is optionally selected from the group consisting of CH₂N(CH₃), CH₂CH₂N(CH₃), and CH₂CH₂N(CH₂CH₃) with the proviso that B is hydrido;

— M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxymethyl, amino, aminomethyl, methylamino, cyano, methyl, trifluoromethyl, methoxy, methylthio, trifluoromethoxy, fluoro, and chloro;

R² is selected from the group consisting of phenyl, 2-thienyl, 2-furyl, 2-pyrrolyl, 2-imidazolyl, 2-thiazolyl, 3-isoxazolyl, 2-pyridyl, and 3-pyridyl, wherein a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to the uracil ring is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹; wherein (a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², and (e) a ring carbon, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹;

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, methyl, ethyl, methoxy, ethoxy, hydroxy, amino, N-methylamino, N,N-dimethylamino, methylthio, trifluoromethyl, pentafluoroethyl,

2,2,2-trifluoroethyl, fluoro, chloro, bromo, amidosulfonyl,

N-methylamidosulfonyl, N,N-dimethylamidosulfonyl, hydroxymethyl,

1-hydroxyethyl, amidocarbonyl, N-methylamidocarbonyl, carboxy, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, amidino, amidocarbonyl, N-methylamidocarbonyl,

N-benzylamidocarbonyl, N-(2-chlorobenzyl)amidocarbonyl,

N-(3-fluorobenzyl)amidocarbonyl, N-(2-trifluoromethylbenzyl)amidocarbonyl, N-(1-phenylethyl)amidocarbonyl, N-(1-methyl-1-phenylethyl)amidocarbonyl, N-(benzylamidosulfonyl, N-(2-chlorobenzyl)amidosulfonyl,

N-ethylamidocarbonyl, N-isopropylamidocarbonyl, N-propylamidocarbonyl, N-isobutylamidocarbonyl, N-(2-butyl)amidocarbonyl,

N-cyclobutylamidocarbonyl, N-cyclopentylamidocarbonyl,

N-cyclohexylamidocarbonyl, guanidino, methyl, ethyl, methoxy, ethoxy, hydroxy, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, carboxy, carboxymethyl, amino, acetamido, trifluoromethyl, pentafluoroethyl,

2,2,2-trifluoroethyl, trifluoroacetamido, aminomethyl, N-methylamino, dimethylamino, methoxyamino, amidosulfonyl, N-methylamidosulfonyl,

N,N-dimethylamidosulfonyl, methanesulfonamido, methoxycarbonyl, fluoro, chloro, bromo, and cyano; <u>and</u>

Y⁰ is selected from the group consisting of:

1-Q^b-4-Q^c-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹benzene,

2-Q^b-5-Q^c-6-R¹⁷-4-R¹⁸-3-R¹⁹pyridine, 2-Q^b-5-Q^c-3-R¹⁶-4-R¹⁷thiophene,

3-Q^b-6-Q^c-2-R¹⁶-5-R¹⁸-4-R¹⁹pyridine, 3-Q^b-5-Q^c-4-R¹⁶-2-R¹⁹thiophene,

3-Q^b-5-Q^c-4-R¹⁶-2-R¹⁹furan, 2-Q^b-5-Q^c-3-R¹⁶-4-R¹⁷furan,

3-Q^b-5-Q^c-4-R¹⁶-2-R¹⁹pyrrole, 2-Q^b-5-Q^c-3-R¹⁶-4-R¹⁷pyrrole,

4-Q^b-2-Q^c-5-R¹⁰thiazole, and 2-Q^b-5-Q^c-4-R¹⁷thiazole;

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, methyl, ethyl, amidino, guanidino, methoxy, hydroxy, amino, aminomethyl,

1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, methylthio, ethylthio, trifluoromethylthio, methylsulfinyl, methylsulfonyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, trifluoromethoxy, fluoro, chloro, hydroxymethyl, carboxy, and cyano[[;]].

Q^b is selected from the group consisting of NR²⁰R²¹, C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴);

—— R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, methyl, and ethyl;

——Q⁶-is-CH₂-

Claim 22 (currently amended): Compound of [[Claim]] <u>claim</u> 21 or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butyl, (R)-2-butyl, (S)-2-butyl, *tert*-butyl, isobutyl, 1-pentyl, 3-pentyl, 2-methylbutyl, 2,2,2-trifluoroethyl, 6-amidocarbonylhexyl, 4-methyl-2-pentyl, 3-hydroxypropyl,

1-methoxy-2-propyl, 2-methoxyethyl, 2-methyl-2-butyl, 3-methyl-2-butyl,

2-dimethylaminopropyl, 2-cyanoethyl, 6-hydroxyhexyl, 2-hydroxyethyl, 2-amidinoethyl, 2-guanidinoethyl, 3-guanidinopropyl, 4-guanidinobutyl,

3-hydroxypropyl, 4-hydroxybutyl, 6-cyanohexyl, 2-dimethylaminoethyl,

3-methylbutyl, 2-methylbutyl, (S)-2-methylbutyl, 3-aminopropyl, 2-hexyl, and 4-aminobutyl;

A is selected from the group consisting of a bond, CH₂, CH₃CH, and CH₂CH₂;

——— M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxymethyl, amino, aminomethyl, cyano, methyl, trifluoromethyl, and fluoro;

R² is selected from the group consisting of

3-amidocarbonyl-5-aminophenyl, 3-amidocarbonyl-5-aminophenyl,

3-amino-5-(N-benzylamidocarbonyl)phenyl,

3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl,

3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl,

```
3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl,
```

3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl,

3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl,

3-amino-5-(N-benzylamidosulfonyl)phenyl,

3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl,

3-amino-5-(N-ethylamidocarbonyl)phenyl,

3-amino-5-(N-isopropylamidocarbonyl)phenyl,

3-amino-5-(N-propylamidocarbonyl)phenyl,

3-amino-5-(N-isobutylamidocarbonyl)phenyl,

3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl,

3-amino-5-(N-cyclobutylamidocarbonyl)phenyl,

3-amino-5-(N-cyclopentylamidocarbonyl)phenyl,

3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 5-amino-2-fluorophenyl,

3-amino-5-hydroxymethylphenyl, 5-amino-3-methoxycarbonylphenyl,

3-amidinophenyl, 3-amino-2-methylphenyl, 5-amino-2-methylthiophenyl,

3-aminophenyl, 3-carboxyphenyl, 3-carboxy-5-aminophenyl,

3-carboxy-5-hydroxyphenyl, 3-carboxymethyl-5-aminophenyl,

3-carboxymethyl-5-hydroxyphenyl, 3-carboxymethylphenyl, 3-chlorophenyl, 2-

chlorophenyl, 3-cyanophenyl, 3,5-diaminophenyl, 3-dimethylaminophenyl,

2-fluorophenyl, 3-fluorophenyl, 2,5-difluorophenyl, 2-hydroxyphenyl,

3-hydroxyphenyl, 3-methanesulfonylaminophenyl, 2-methoxyphenyl,

3-methoxyphenyl, 3-methoxyaminophenyl, 3-methoxycarbonylphenyl,

2-methylaminophenyl, 3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, 4-

methylphenyl, phenyl, 3-trifluoroacetamidophenyl, 3-trifluoromethylphenyl, and 2-

trifluoromethylphenyl, 5-amino-2-thienyl, 5-amino-3-thienyl,

3-bromo-2-thienyl, 3-pyridyl, 4-pyridyl, 2-thienyl, and 3-thienyl;

Y⁰ is selected from the group consisting of:

1-Q⁵-4-Q⁵-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹benzene.

2-Q^b-5-Q^c-6-R¹⁷-4-R¹⁸-3-R¹⁹pyridine,

3-Q^b-6-Q^c-2-R¹⁶-5-R¹⁸-4-R¹⁹pyridine.

3-Q^b-5-Q^c-4-R¹⁶-2-R¹⁹thiophene, and 2-Q^b-5-Q^c-3-R¹⁶-4-R¹⁷thiophene;

——Q⁶ is CH₂.

R¹⁶ and R¹⁹ are independently selected from the group consisting of hydrido, amidino, amino, aminomethyl, methoxy, methylamino, hydroxy, hydroxymethyl, fluoro, chloro, and cyano; <u>and</u>

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrido, fluoro, chloro, hydroxy, hydroxymethyl, amino, carboxy, and cyano[[;]].

Q^b is C(NR²⁵)NR²³R²⁴;

R²³, R²⁴, and R²⁵ are independently hydride or methyl;

Claim 23 (currently amended): Compound of [[Claim]] <u>claim</u> 22 or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butyl, (R)-2-butyl, (S)-2-butyl, *tert*-butyl, isobutyl, 1-pentyl, 3-pentyl, 2-methylbutyl, 2,2,2-trifluoroethyl, 6-amidocarbonylhexyl, 4-methyl-2-pentyl, 3-hydroxypropyl, 1-methoxy-2-propyl, 2-methoxyethyl, 2-methyl-2-butyl, 3-methyl-2-butyl, 2-dimethylaminopropyl, 2-cyanoethyl, 6-hydroxyhexyl, 2-hydroxyethyl, 2-amidinoethyl, 2-guanidinoethyl, 3-guanidinopropyl, 4-guanidinobutyl, 3-hydroxypropyl, 4-hydroxybutyl, 6-cyanohexyl, 2-dimethylaminoethyl, 3-methylbutyl, 2-methylbutyl, (S)-2-methylbutyl, 3-aminopropyl, 2-hexyl, and 4-aminobutyl;

A is selected from the group consisting of a bond, CH₂, CH₃CH, and CH₂CH₂;

—— M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxymethyl, amino, aminomethyl, cyano, methyl, trifluoromethyl, and fluoro;

R² is selected from the group consisting of
3-amidocarbonyl-5-aminophenyl, 3-amino-5-(N-benzylamidocarbonyl)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl,
3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl,
3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl,
3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl,

- 3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl,
- 3-amino-5-(N-benzylamidosulfonyl)phenyl,
- 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl,
- 3-amino-5-(N-ethylamidocarbonyl)phenyl,
- 3-amino-5-(N-isopropylamidocarbonyl)phenyl,
- 3-amino-5-(N-propylamidocarbonyl)phenyl,
- 3-amino-5-(N-isobutylamidocarbonyl)phenyl,
- 3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl,
- 3-amino-5-(N-cyclobutylamidocarbonyl)phenyl,
- 3-amino-5-(N-cyclopentylamidocarbonyl)phenyl,
- 3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 3-aminophenyl,
- 3-carboxy-5-aminophenyl, 3-chlorophenyl, 3,5-diaminophenyl,
- 3-dimethylaminophenyl, 3-hydroxyphenyl, 3-methanesulfonylaminophenyl,
- 3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, phenyl, and
- 3-trifluoroacetamidophenyl, 3-bromo-2-thienyl, 2-thienyl, and 3-thienyl; and
- Y⁰ is selected from the group consisting of 5-amidino-2-thienylmethyl, 4-amidinobenzyl, 2-fluoro-4-amidinobenzyl, and 3-fluoro-4-amidinobenzyl.

Claim 24 (currently amended): Compound of [[Claim]] claim 17, where said compound is selected from the group of the Formula:

$$\begin{array}{c|c} & & & \\ & & \\ & & & \\ & & & \\ & & & \\ & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ &$$

or a pharmaceutically acceptable salt thereof, wherein;

 R^2 is 3-aminophenyl, B is 2,2,2-trifluoroethyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is (S)-2-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 5-amino-2-fluorophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 2-methyl-3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is ethyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is ethyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-propenyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

R² is 3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

 R^2 is 3-aminophenyl, B is (R)-2-butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-propynyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 3-pentyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is hydrido, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is ethyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-methypropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-propyl, A is CH₃CH, Y⁰ is 4-amidinobenzyl, and M is CH;

 R^2 is 3-aminophenyl, B is propyl, A is single bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is CH;

 R^2 is 3-aminophenyl, B is 6-amidocarbonylhexyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is tert-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is tert-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 3-hydroxypropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-methylpropyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

 R^2 is 3-aminophenyl, B is butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;


R² is 3-aminophenyl, B is 1-methoxy-2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-methoxyethyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

is N;

R2 is 3-aminophenyl, B is 2-propyl, A is single bond, Y⁰ is 5-amidino-2thienylmethyl, and M is CH; R² is 3-aminophenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidino-3fluorobenzyl, and M is CH; R² is 3-carboxyphenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH; R² is 3-aminophenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidino-3fluorobenzyl, and M is CH; R2 is 3-aminophenyl. B is 2,2,2-trifluoroethyl, A is single bond, Y⁰ is 4amidinobenzyl, and M is N; R2-is-3-aminophenyl, B is (S)-2-butyl, A is single bond, Y⁰-is-4-amidinobenzyl, and M is N: R2 is 5-amino-2-fluorophenyl, B is isopropyl, A is single bond, Y⁰ is 4amidinobenzyl, and M is N; R2 is 2-methyl-3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4amidinobenzyl, and M is N; R2-is 3-aminophenyl, B is ethyl, A is single bond, Y⁰-is 4-amidinobenzyl, and M is ₩; R2 is 3-aminophenyl, B is ethyl, A is single bond, Y0 is 4-amidino-2-fluorobenzyl, and M is N; R2 is 3-aminophenyl, B is 2-propenyl, A is single bond, Y is 4-amidinobenzyl, and M is N: R2 is 3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidino-2fluorobenzyl, and M is N: R2 is 3-aminophenyl, B is isopropyl, A is single bond, Y is 4-amidinobenzyl, and M-is-N;

R2 is 3-aminophenyl, B is 2-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M



- -----R2 is 3-aminophenyl, B is 2-methoxyethyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-aminophenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidino-3-fluorobenzyl, and M is N;
- ———R2-is 3-carboxyphenyl, B is 2-propyl, A is-single bond, Y⁰ is 4-amidinobenzyl, and M is-N;
- R² is 3-aminophenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidino-3-fluorobenzyl, and M is CH;
- R² is 3-amino-5-carboxyphenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R^2 is 3-amino-5-carbomethoxyphenyl, B is isopropyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;
- R² is 3-aminophenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CCI;
- R^2 is 3-amino-5-carboxamidophenyl, B is isopropyl, A is a bond, \dot{Y}^0 is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-benzyl-N-methylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(2-phenyl-2-propyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(2,4-dichlorobenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(4-bromobenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;

- R^2 is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R^2 is 3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(3-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R^2 is 3-amino-5-(N-isobutylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;
- R^2 is 3-amino-5-(N-cyclobutylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;
- R^2 is 3-amino-5-(N-cyclopentylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-cycloheptylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(2-pyridylmethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(3-pyridylmethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(2-(4-methoxyphenyl)ethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(3-phenylpropyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(2,2-diphenylethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(2-naphthylmethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R² is 3-amino-5-(N-(1,2,3,4-tetrahydronaphth-2-ylmethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-propyl, A is a bond, Y⁰ is 4-amidino-3-fluorobenzyl, and M is CH;

R² is 3,5-diaminophenyl, B is 2,2,2-trifluoroethyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;

 R^2 is 3,5-diaminophenyl, B is (S)-2-butyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;

R² is 3,5-diaminophenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3,5-diaminophenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidino-2-fluorobenzylbenzyl, and M is CH;

 R^2 is 3,5-diaminophenyl, B is ethyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;

R² is 3,5-diaminophenyl, B is ethyl, A is a bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

 R^2 is 3-amino-5-carboxyphenyl, B is 2,2,2-trifluoroethyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-carboxyphenyl, B is (S)-2-butyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-carboxyphenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidino-2-fluorobenzylbenzyl, and M is CH;

R² is 3-amino-5-carboxyphenyl, B is ethyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-carboxyphenyl, B is ethyl, A is a bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;


R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is 2,2,2-trifluoroethyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is (S)-2-butyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;

 R^2 is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y^0 is 4-amidino-2-fluorobenzylbenzyl, and M is CH;

- R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is ethyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is CH;
- R^2 is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is ethyl, A is a bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is CH; <u>or</u>
- R2 is 3-amino-5-carboxyphenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-carbomethoxyphenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-carboxamidophenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-benzyl-N-methylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(2-phenyl-2-propyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(2,4-dichlorobenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(4-bromobenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B-is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;

- R2 is 3-amino-5-(N-(3-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-isobutylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-cyclobutylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2-is-3-amino-5-(N-cyclopentylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰-is-4-amidinobenzyl, and M is N;
- R^2 is 3-amino-5-(N-cycloheptylamidocarbonyl)phenyl, B is isopropyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is N;
- R2 is 3 amino 5 (N-(2-pyridylmethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2-is 3-amino-5-(N-(3-pyridylmethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(2-(4-methoxyphenyl)ethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(3-phenylpropyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(2,2-diphenylethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(2-naphthylmethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R2 is 3-amino-5-(N-(1,2,3,4-tetrahydronaphth-2-ylmethyl)amidocarbonyl)phenyl, B is isopropyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;
- R^2 is 3-carboxyphenyl, B is 2-propyl, A is a bond, Y^0 is 4-amidinobenzyl, and M is CCI[[;]].
- R2 is 3-aminophenyl, B is 2-propyl, A is a bond, Y⁰ is 4-amidino-3-fluorobenzyl, and M is N;



——R2-is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is ethyl, A is a bond, Y⁰ is 4-amidinobenzyl, and M is N;

R2-is-3-amino-5-(N-benzylamidocarbonyl)phenyl, B is ethyl, A is a bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is N.

25-37 (cancelled)

38 (currently amended) A composition for inhibiting thrombotic conditions in blood comprising a compound of any one of Claims 8, 16, 24, 32, and 37 claims 1, 2, and 17-24 and a pharmaceutically acceptable carrier.

Claim 39 (cancelled)

Claim 40 (currently amended): A method for inhibiting thrombotic conditions in blood comprising adding to blood a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 41 (currently amended): A method for inhibiting formation of blood platelet aggregates in blood comprising adding to blood a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 42 (currently amended): A method for inhibiting thrombus formation in blood comprising adding to blood a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 43 (currently amended): A method for treating or preventing venuous thromboembolism and pulmonary embolism in a mammal comprising administering to the mammal a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 44 (currently amended): A method for treating or preventing deep vein thrombosis in a mammal comprising administering to the mammal a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 45 (currently amended): A method for treating or preventing cardiogenic thromboembolism in a mammal comprising administering to the mammal a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 46 (currently amended): A method for treating or preventing thromboembolic stroke in humans and other mammals comprising administering to the mammal a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 47 (currently amended): A method for treating or preventing thrombosis associated with cancer and cancer chemotherapy in humans and other mammals comprising administering to the mammal a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 48 (currently amended): A method for treating or preventing unstable angina in humans and other mammals comprising administering to the mammal a therapeutically effective amount of a composition of any one of Claims 38 and 39 claim 38.

Claim 49 (currently amended): A method for inhibiting thrombus formation in blood comprising adding to blood a therapeutically effective amount of a compound of any one of Claims 1 through 37 claims 1, 2, and 17-24 with a therapeutically effective amount of fibrinogen receptor antagonist.

Claim 50 (cancelled)